SPANAWAY


PIERCE County

Lake ID: SPAPI1
Ecoregion: 2

Lake Spanaway is located ten miles south of Tacoma, and 0.5 mile west of Spanaway. It is fed by drainage from a swampy area, and drains via Spanaway Creek to Clover Creek and Lake Steilacoom. Daron Island lies in the north portion of the lake

Area (acres)	Maximum Depth (ft)
280	28
Volume (ac-ft)	Shoreline (miles)
4600	4.36

Mean Depth (ft)	Drainage (sq mi)			
16	17			
Altitude (ft abv msl)	Latitude	Longitude		
320	47 07 11.	122 26 45.		



Primary Station	Station # 1	latitude: 47 06 47.0	longitude: 122 27 01.7
	Description:	In deep part of lake, due west of pushore of island	ablic boat launch and south of eastern
Secondary Station	Station # 2	latitude: 47 06 24.4	longitude: 122 26 58.7
	Description:	2: Across from boat launch, approximately 500 feet east of west shore at point directly across from boat launch	
Secondary Station	Station # 3	latitude:	longitude:
	Description:	In horizontal middle of lake near south end, directly south of boat launc and directly east of a small cove just north of southern portion of lake; no coordinates recorded	

Trophic State Assessment	for	1998			SPANAWAY
Analyst: KIRK SMITH			TSI_Secchi:	44	
			TSI_Phos:	48	
			TSI_Chl:	54	
			Narrative TSI: ^a	ME	

Spanaway Lake is an urban lake bordered by a large city park along its northeast shore. The lake has suffered numerous blue-green algal blooms in recent history and shows a tendency to go anoxic in the hypolimnion during the summer, resulting in periodic internal nutrient release and very high hypolimnetic phosphorus concentrations. Records show that the lake was once dredged, which may have provided for many years of artificially clear water. Spanaway Lake is fed by a stream originating from a wetland. The wetland, park, and golf course are potential sources of nutrients to the lake. Natural shoreline is altered in the park and fertilizer runoff from the golf course may contribute to the eutrophication--although most of the runoff likely enters near the outlet of the lake. The user survey suggests the water clarity is deteriorating through the years with only one respondent out of 11 believing the clarity has improved. Detractors to the use of the lake, according to survey respondents, include the poor water clarity and the abundant Canada geese. All respondents said they'd rather have clearer water than fewer aquatic plants. A significant Aphanizomenon bloom was observed in August.

Although the lake has been productive for largemouth bass fishing in recent history, it's questionable whether the extensive anoxia in the summertime hypolimnion is conducive to a good trout fishery. Having both a good trout and bass fishery is important to the residents who responded to the survey. Water clarity may impair primary contact recreation at times but the overall Secchi TSI suggests more mesotrophic conditions. We assigned an overall assessment of meso-eutrophic. The habitat survey revealed significant human influence (lawns, buildings and docks) on the habitat of the lake. There was also a high population of geese on the lake to take advantage of all the human influences conveniently provided.

We recommend that a nutrient criterion be set at the ecoregion action value for lower mesotrophic Puget Lowland lakes, 20 ug/L.

^a E=eutrophic, ME=mesoeutrophic, M=mesotrophic, OM=oligomesotrophic, O=oligotrophic

Chemis	stry l	<u>Data</u>							SI	PANAWAY
Date	Time	Strata		Tot N (mg/L)	TN:TP	Chloro- phyll (ug/L)	Fecal Col. Bacteria (#/100mL)	Hardness (mg/L)	Calcium (ug/L)	Turbidity (NTU)
Station 0										
8/19/1998		L					8			
		L					13			
9/23/1998		L					8			
		L					12			
Station 1										
6/12/1998		E	12.8	.818	64	11		41.7	10800	1.6 J
		Н	29.3	1.11	38					
7/30/1998		E	15.8	.476	30	6.8				1.5 J
		Н	340	2.31	7					
8/19/1998		E	16.6	.4	24	15.5				1.5
		Н	15.3	1.63	107					
9/23/1998		Е	21.3	.356	17	11.2				1.5
		Н	318	2.39	8					
Station 3										
6/12/1998		E	12.7			8.8				
9/23/1998		E	29.2	.339	12					

Strata: L=lake surface, E=epilimnion, H=hypolimnion; Qualifier: J=Estimate, U=Less than

Watershed Survey

SPANAWAY

Land Uses (1 = Primary, 2 = Secondary, etc.)	Survey Date: 9/23/1998
Agriculture(commercial, not hobby) 4 Commercial, Industrial	1 Residential 2 Park, forest or natural
3 Major transportation	

Impervious surfaces (Roads and parking area): Partially Curbed

Observations (check mark denotes presence)

BMP's			
Natural vegetation lacking along shoreline.			
Odors			
Cattle			
Fertilizers and weed killers appear to be used in residential or agriculture area	✓		
Yes, manicured lawns and a public golf course.			
Buffer zones around streams and wetlands			
Irrigation □			
		Survey Id:	30

Habitat Survey Summary Report

SPA	NAWAY
Date of Visit:	8/19/1998

•			SPA	NAWAI
Data are averages of 10 Sta	tions Surveyed		Date of Visit:	8/19/199
Vegetation Type (Avg. o	only of sites w/	vegetation present; 1=conifere	ous, 3=decidu	ious)
Canopy Layer Avg:	1.6	Number of stations with canopy:	10	
Understory Avg:	2.7	Number of stations with understory:	10	
Percent Areal Coverage	(0 = absent, 1)	=<10%, 2 = 10-40%, 3 = 40-75%, 4	=>75%)	
Canopy Layer:	trees > 0.3 m DBI	H	1.5	
	trees< 0.3 m DBH	I	0.6	
Understory:	woody shrubs saj	plings	2.2	
	tall herbs, forbs	grasses	1.5	
Ground Cover:	woody shrubs see	edlings	2.1	
	herbs, forbs, gras	sses	2.7	
	standing water or	inundated veg	0.9	
	barren or buildin	gs	1.3	
Substrate Type	bedrock		0.0	
(within	boulders		0.1	
shoreline plot):	cobble/gravel		1.8	
	loose sand		0.7	
	other fine soil/sed	iment	0.6	
	vegetated		2.5	
	other		0.8	
Bank Features:	angle (O:<30; 1: 3	30-75; 2:nr vertical)	0.5	
	vertical dist (M fr	om wtrln to high wt):	0.2	

Human Influence	(0 = absent, 1 = adjacent to or behind	plot, 2 = present within plot)
	buildings	1.0
	commercial	0.0
	park facilities	0.4
	docks/boats	1.2
	walls, dikes, or revetments	0.9
	litter, trash dump, or landfill	0.2
	roads or railroad	0.0
	row crops	0.0
	pasture or hayfield	0.0
	orchard	0.0
	lawn	1.4
	other	0.2
Physical Habitat Cha	racteristics	
	station depth (at 10 m from shore)	1.8
Bottom Substrate (0 =	= absent, 1 = <10%, 2 = 10-40%, 3 = 4	40-75%, 4 = >75%
	bedrock	0.0
	boulders	0.1
	cobble	1.6
	gravel	1.9
	sand	1.2
	silt	1.2
	woody debris	0.3
Macrophyte Areal Co	overage (0 = absent, 1 = <10%, 2 = 10	-40%, $3 = 40-75%$, $4 = >7$
	submergent	0.5
	emergent	1.2
	floating	0.3
	total weed cover	1.1
Do macrophytes ex	xtend lakeward (-1 = yes, 0 = no)	-0.3
Fish Cover (0 = abser	nt, 1 = Present but sparse, 2 = modera	ate to heavy)
	aquatic weeds	0.8
	snags	0.0
	brush or woody debris	0.9
	inundated live trees	0.1
	overhanging vegetation	2.0
	rock ledges or sharp dropoffs	0.0
	boulders	0.0
	human structures	1.2

Questionnaire SPANAWAY

Questionnaire						SPANAV	VA'
Results compiled from	11 Surveys.		Average time (years) respo	ndents spent o	n lake:	10.2
Did the following add (+1), detract (-1),	or have no effect (0) on your enjoym	ent of the lak	e today?		
Types of WaterCraft:	-0.5 V	View:	0.7	Γ	Distance to Lake	e:	0.5
Public Access:	-0.3	Swim Beach:	-0.5	C	Canada Geese:		-0.9
Water Clarity:	-0.7 V	Water Qual. for Sw	im: -0.6				
Fishing Quality:	0.0	Aquatic Plants:	-0.3				
On a scale of 1 (poor) to	5 (excellent), ho	ow would you rate	water quality toda	y? 2.0			
Which would you rathe	er have, 1 or 2?						
1) Better fishing and mo	ore natural habitat	, or 2) clearer water	r?	1.6			
1) Better fishing and mo	ore natural habitat	, or 2) fewer aquati	c plants?	1.5			
1) Clearer water, or 2) f	ewer aquatic plan	ts?		1.0			
How important is each			ou (1 = very under	sirabla 5= va	ry desirable):		
-	3.6		•		,	4.4	
Restricted Watercraft: Plant Growth:	2.3	Good Warmwtr Good Swimmin	i ioming.		tural Scenery: blic Beach:	2.2	
Natural Shoreline:	3.5	Less Algae:	g. 4.1 4.2		nada Geese:	1.6	
No Odors:	4.3	Public Access:	2.5	Ca	mada Geese.	1.0	
Good Coldwtr Fishing:	4.4	Clear Water:	4.1				
-							
Tabulated Results				χ.	v d		
Survey		Rent or	Primary	Purchase	Vater Clarity- Has it		
ID Date	Residency	Own	Activity*	Factor?	Changed?	When?	
53 7/22/1998 Reside	nt Permanent	Rent	10	✓	Worse	1991	
54 7/25/1998 Reside	nt Permanent	Rent	8	✓	Worse	1989-19	990
55 8/10/1998 Reside	nt Permanent	Rent	10		Worse	1994	
56 8/16/1998 Reside	nt Permanent	Rent	8	✓	Worse	1996	
57 8/25/1998 Reside	nt Permanent	Rent	3	✓	Worse	1997	
58 7/29/1998 Reside	nt Permanent	Rent	1	✓	Worse		
69 8/24/1998 Reside	nt Permanent	Rent	7		No		
73 8/31/1998 Reside	nt Permanent	Rent	several of the abo	ove \square	No		
75 12/31/1998 Residen	nt	Rent			Unknow	n	
76 8/29/1998 Visitor			2		Better	1993	
83 8/19/1998 Visitor			10		Unknow	n	

^{* 1=}canoe/kayak, 2=fish, 3=pers. wtrcrft, 4=mtrboat, 5=sail, 6=swim/wade, 7=watch wldlf, 8=ski, 9=windsurf, 10=relaxing

Zooplankton Report


SPAPI1

Date 6/12/1998

Station: 1 Sample ID 14 Too few to count

Number of organisms measured: 14

Group	Percent	Group Percent
Cladoceran	21.4%	Small < 1mm 85.7%
Copepod	78.6%	Large >= 1mm 14.3%
Other		Ratio of large to Small: 0.17
		Average size (mm): 0.51



Date	Time	Temp- erature (F)	Secchi (ft)	Color (1-greens, 11-browns	Bright- ness (pct)	(1-none,	Rainfall (0-none, 5-heavy)	Aesthetics (1-bad, 5- good)	Swimming (1-poor, 5- good)	Geese (#)	Waterfowl (besides geese #)	Boats- Fishing (#)	Boats- Skiing (#)
Station 1													
5/16/1998	Sample	13.3 er: THOMP	11 SON	6 Remark	100 s:	4	4	4	4	5	6	1	0
5/30/1998	Sample	15.6 er: THOMP	10.25 SON	6 Remark	100 s:	2	2	3	3	8	6	6	1
6/12/1998		18.8889	10.5	6	30			3	3	0	0	0	0
	Sample	er: SMITH		Remark		DOR AT 7 ME ibration failing			T SITE #3. The C)xygen rest	alt is qualified as	an estimate du	e to
6/13/1998	Sample	18.9 er: THOMP	10 SON	6 Remark	100 s:	2	1	3	3	20	6	5	2
6/28/1998		19.4	7.5	6	0	3	1	3	3	12	6	0	4
	Sample	er: THOMP	SON	Remark	s: ALOT ALGAI								
7/11/1998		20.6	12	6	75	2	3	3	3	26	2	12	1
	Sample	er: THOMP	SON	Remark		17TH - WARN 'N RISK".	ING POSTED	AT COUNTY	PARK ON LAKE	E - "HIGH	LEVELS OF FEC	CAL COLIFOR	M; SWIM
7/26/1998		25.6	9	6	0	2	1	1	1	28	0	2	23
	Sample	er: THOMP	SON	Remark		STILL CLOSE JULY 17, 199		MERS DUE TO	HIGH LEVELS	OF FECAI	COLIFORM. L	AKE HAS BE	EN CLOSED
7/30/1998			10.89	6	100	2		1	1	74	110	0	0
	Sample	er: SMITH		Remark		N THE WATE APPLICATION	,		NG. PARK ATTE	ENDENT S	SAID THERE MA	Y HAVE BEI	EN ILLEGAL
8/10/1998		24.4	9.5	6	0	2	1	1	1	25	8	5	5
	Sample	er: THOMP	SON	Remark				ED BY TACOM E JULY 17, 199		PT. DUE T	O HIGH LEVEL	S OF FECAL (COLIFORM.
8/19/1998			8.58	3	0			3	3	0	0	0	0
	Sample	er: SMITH		Remark		NIZOMENON			STRONG H2S C NO BOATS OR		OM 7.5M SAMPL FOWL	.E. ABUNDA	NT

Date	Time	Temp- erature (F)	Secchi (ft)	Color (1-greens, 11-browns	Bright- ness (pct)	Wind (1-none, 5-gusty)			Swimming (1-poor, 5- good)	Geese (#)	Waterfowl (besides geese #)	Boats- Fishing (#)	Boats Skiing (#)
8/23/1998		21.1	8.5	6	25	2	1	1	1	44	8	5	1
	Sampler: THOMPSON			Remark	colifo		D PER HEAI	TH DEPT. DUE	E TO FECAL				
9/6/1998		22.2	15.5	6	0	1	1	3	3	24	6	3	7
	Sample	r: THOMP	SON	Remark		H DEPT. REC			E ON 9/14/98. G	REAT CL	ARITY TODAY I	BUT MORE W	EEDS
9/19/1998		20	9.25	6	25	1	4	3	3	16	6	3	3
	Sample	г: ТНОМР	SON	Remark	es: LAKE I 9/4/98.	REOPENED							
9/23/1998			10.73	3	0			4	3	30	260	2	0
	Sample	r: SMITH		Remark		EAGLE SPOT G H2S IN HY		AT PLATFORM	M ON SOUTH EÌ	ND; FEC#	2 AT PLATFORN	AT SWIM B	EACH.
9/27/1998		16.7	10.5	6	100	2	5	3	3	8	8	5	0
	Sampler: THOMPSON			Remark	s:								
10/18/1998		14.4	8.5	8	50	1	3	1	1	6	400	7	1
	Sample	r: THOMP	SON	Remark		GREEN ALGA OTS ON THE		TE ALONG SH	ORE. LAKE WA	ATER HAS	TURNED BROY	WN! HUNDR	EDS(400+